



## Suicides in the Military: The Post-Modern Combat Veteran and the Hemingway Effect

<b>AUTHORS:</b>	Carl Andrew Castro, Ph.D. Sara Kintzle, M.S.W., Ph.D.
<b>DATE:</b>	2014
<b>PUBLICATION:</b>	<i>Current Psychiatry Reports</i> , 16(8), 460-469
<b>PUBLICATION TYPE:</b>	Peer-Reviewed Journal Article
<b>KEYWORDS:</b>	Military, suicide, veteran, combat, interpersonal -psychological theory of suicide, military transition theory, modern veteran, Hemingway effect

### ABSTRACT

“Suicides in the military have increased over the last ten years. Much effort has been focused on suicide prevention and treatment, as well as understanding the reasons for the sharp increase in military suicides. Despite this effort, the definitive causes of military suicides remain elusive. Further, highly effective suicide prevention and treatment approaches have not yet been developed. The purpose of this article is to present a short review of the current state of suicide prevention interventions within the context of the military. The root causes of suicidal behavior and the role of combat in the military are each discussed. Interpersonal-psychological theory of suicide and the military transition theory are introduced as guiding frameworks for understanding suicides and suicidal behavior amongst active military personnel and military veterans. The article concludes with a set of recommendations for moving forward in understanding and addressing suicides in the military.”

### RESEARCH HIGHLIGHTS

- Between 1990 and 2007, the military suicide rate was below the civilian suicide rate. However, in recent years, the military suicide rate has increased. In a 2012 data report, the VA reported that an active duty service member dies by suicide every 36 hours and more than 8,000 military veterans died by suicide each year. In an effort to address suicide in military related individuals, Castro and Kintzle review the current state of suicide prevention interventions within the context of the military.
- Previous researchers have found that combat is associated with an increased rate of PTSD and depression. However, researchers have not established a direct link between combat and suicide, thus military suicides is evaluated in context with an increase in mental health issues of veterans and service members.
- Castro and Kintzle draw connections between the life of Ernest Hemingway, who died by suicide, and aging veterans who died by suicide. Discussing this in line with interpersonal-psychological theory of suicide, Castro and Kintzle show that as some veterans age their perceived burdensomeness and lack of belonging increases. The increase of burdensomeness and lack of belongingness is coupled with their acquired capability of knowing how to use weapons from their service which could result in some veterans turning towards suicide.
- Addressing suicide by targeting sense of burden and sense of not belonging could help family members, friends, and medical providers prevent aging veterans and service members from dying by suicide.

# BRIEF

## FOR PRACTICE

Practitioners should monitor the wellness of their aging veterans, anticipating the needs of elder veterans to reduce depression and PTSD symptoms and suicide. Practitioners should also assist their elder veterans with obtaining necessary support, which might include recommending individual or group counseling sessions. Practitioners should communicate with loved ones to ensure the aging veteran has a support system that minimizes feelings of not belonging and burdensomeness. Practitioners should consider discussing with loved ones ways to keep the elderly veteran involved in family decisions and activities. The interpersonal-psychological theory of suicide used in this study posits that individuals who die by suicide must have an acquired capability, or be familiar with using weapons. Recent CDC reports on civilian suicides show that older men who die by suicide often by means of a gun or similar weapon. Thus, practitioners and family members with an aging veteran, especially a male, should discuss with their veteran ways to feel fulfilled and engaged. While serving, many service members share a sense of comradery and pride that is absent among civilians.

Noted from the service should continue this comradery by participating in social, community, and service events with other veterans. Veterans struggling with depression or PTSD symptoms or feeling like they do not belong should discuss the symptoms and feelings with their physician. To increase one's sense of belonging and make a smoother transition, veterans should establish and re-establish relationships and an identity that is not strongly tied with their military experience.

## FOR POLICY

The DoD might develop initiatives and interventions aimed at addressing the current physical and psychological needs, in addition to ensuring feelings of belongingness within are maintained for active-duty service members. The VA might implement similar initiatives and interventions as the DoD for aging veterans, being proactive rather than reactive.

## ABSTRACT

Older veterans before a need arises. Additionally, since 45-64 and aged 85 and older in the United States have the highest and depression. How “Suicides in the military have increased over the last highest suicide rates, respectively, policymakers might established initiatives to prevent and treat suicide in military populations. Unlike veterans of military prevention and treatment as well as understanding half of the US population with an increasing number of military suicides. To increase feelings of belongingness, the VA might consider highly effective support groups to reduce perceived burden and lack of belongingness. The VA have made many improvements to help veterans with PTSD and depressive symptoms, which might indirectly reduce suicide risk and attempts. To reduce the stigma and barriers associated with asking for help, the DoD might work to prevent mental illness and suicide from the first day of service through resilience skill training and creating a more supportive culture of report and understanding between leaders and colleagues. Furthermore, the VA might monitor their well-being for 2-3 years after service to ensure veterans are transitioning well and not suffering. Interpersonal-psychological theory of suicide and the military transition theory are introduced as guiding frameworks for understanding suicides and best active military personnel and military veterans. The identification and understanding of mechanisms that reduce perceived burden and lack of belongingness and conditions for providing forward it may also be beneficial to closely investigate the type of civilian communities and organizations (for example, Team Red, White, and Blue, and the Team RUDicom) that supposedly produce the most useful support for veterans exhibiting PTSD or depressive symptoms. Future researchers should also continue to evaluate the effectiveness of various preventative mental health interventions and treatments in reducing suicide risk.

## RESEARCH HIGHLIGHTS

- Between 1990 and 2007, the military suicide rate was below the civilian suicide rate. However, in recent years, the military suicide rate has increased. In a 2012 data report, the VA reported that an active duty service member dies by suicide every 36 hours and more than 8,000 military veterans died by suicide each year. In an effort to address suicide in military related individuals, Castro and Kintzle reviewed the current research and depression. Previous researchers is associated with an established initiative to prevent and treat suicide in military populations. Unlike veterans of military context with an increasing number of military suicides. To increase feelings of belongingness, the VA might consider highly effective support groups to reduce perceived burden and lack of belongingness. The VA have made many improvements to help veterans with PTSD and depressive symptoms, which might indirectly reduce suicide risk and attempts. To reduce the stigma and barriers associated with asking for help, the DoD might work to prevent mental illness and suicide from the first day of service through resilience skill training and creating a more supportive culture of report and understanding between leaders and colleagues. Furthermore, the VA might monitor their well-being for 2-3 years after service to ensure veterans are transitioning well and not suffering. Interpersonal-psychological theory of suicide and the military transition theory are introduced as guiding frameworks for understanding suicides and best active military personnel and military veterans. The identification and understanding of mechanisms that reduce perceived burden and lack of belongingness and conditions for providing forward it may also be beneficial to closely investigate the type of civilian communities and organizations (for example, Team Red, White, and Blue, and the Team RUDicom) that supposedly produce the most useful support for veterans exhibiting PTSD or depressive symptoms. Future researchers should also continue to evaluate the effectiveness of various preventative mental health interventions and treatments in reducing suicide risk.

## AUTHOR INFORMATION

**Carl Andrew Castro, Ph.D.<sup>1,2</sup>**  
**Sara Kintzle, M.S.W., Ph.D.<sup>2</sup>**

---

<sup>1</sup> carl.castro@usc.edu  
<sup>2</sup> School of Social Work, Center for Innovation and Research on Veterans and Military Families, University of Southern California